



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : <b>A61K 9/12</b>	<b>A1</b>	(11) International Publication Number: <b>WO 00/66084</b> (43) International Publication Date: 9 November 2000 (09.11.00)
<p>(21) International Application Number: PCT/US00/12092</p> <p>(22) International Filing Date: 3 May 2000 (03.05.00)</p> <p>(30) Priority Data: 60/132,472 4 May 1999 (04.05.99) US Not furnished 2 May 2000 (02.05.00) US</p> <p>(71) Applicant (for all designated States except US): ARADIGM CORPORATION [US/US]; 3929 Point Eden Way, Hayward, CA 94545 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): RUBSAMEN, Reid, M. [US/US]; 1500 Serafin Road, Alamo, CA 94507 (US); COLE, Robert [US/US]; Hayward, CA (US).</p> <p>(74) Agent: BOZICEVIC, Karl; Bozicevic, Field &amp; Francis LLP, Suite 200, 200 Middlefield Road, Menlo Park, CA 94025 (US).</p>	<p>(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>	
(54) Title: AEROSOL FORMULATIONS AND DEVICES FOR INCREASING LIBIDO IN WOMEN VIA ACUTE TESTOSTERONE ADMINISTRATION		
<p>(57) Abstract</p> <p>The libido of adult human female patients is increased by the intrapulmonary delivery of testosterone. A formulation of testosterone is aerosolized and inhaled into a patient's lungs where particles of testosterone deposits on lung tissue and then enter the patient's circulatory system. The patient's testosterone level is enhanced well above baseline levels for a short period and subsides to baseline levels with normal metabolism thereby providing desired short term effects on enhanced libido without undesirable effects of long term enhanced testosterone levels. Additional formulations are provided including formulations for aerosolized delivery of sildenafil citrate which are delivered to male or female patients.</p>		

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CN	Cameroon	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

AEROSOL FORMULATIONS AND DEVICES FOR  
INCREASING LIBIDO IN WOMEN VIA  
ACUTE TESTOSTERONE ADMINISTRATION

5

FIELD OF THE INVENTION

This invention relates generally to aerosol formulations and devices for treating women with a decreased libido. More specifically, the invention relates to acute non-invasive administration of testosterone by inhalation to enhance libido over a discrete period of time.

10

BACKGROUND OF THE INVENTION

The presence of a normal amount of libido, defined as the urge to engage in sexual activity, is an important component of an individual's well-being. In both men and women the primary naturally occurring hormone responsible for libido is testosterone. In males, the baseline testosterone level is a relatively constant throughout life, decreasing slowly in old age. In contrast, women elaborate testosterone only as part of the process of ovulation. Each maturing follicle produces testosterone at the mid-point of the menstrual cycle, consistent with observations that female libido peaks with ovulation. As a woman ages, the number of maturing follicles per month decreases, and there is a decreasing total amount of testosterone produced.

20

A common complaint of post menopausal women is decreased libido. This decrease in libido is characterized by a lack of interest in sexual intercourse, the lack of ability to achieve orgasm, or decrease in intensity of orgasm. It is important to note that this decrease in libido is often associated with a profound sense of loss of a once normal and active interest in sexual activity.

25

Clinicians frequently confronted with the problem of managing female patients presenting with decreased libido have limited tools to address the problem. Testosterone is available as an oral preparation and can be given, for instance, in combination with estrogen to restore testosterone levels. However, the replacement of the once pulsatile endogenous delivery of testosterone with the sustained blood level of the hormone produces unwanted side effects. Women taking testosterone for a few weeks typically begin to complain of the emergence of secondary sexual characteristics such as unwanted body hair, oily hair, and,

30

with prolonged a use, deepening voice. For this reason, oral testosterone replacement therapy is not a practical solution for most patients with decreased libido.

Other forms of testosterone replacement therapy for women are being explored. A transdermal patch capable of delivering a steady rate of testosterone is being tested for use in women. As with oral testosterone replacement therapy, the study state blood levels of testosterone produced via transdermal delivery are likely to be associated with the same side effect profile issues.

It is recognized that testosterone in females decreases with age (Human Biology, May 1980, volume of 52, No. 2, pages 181-0191. ). It is also known that sexual motivation in post menopausal women is associated with the levels of exogenously introduced testosterone (Psychosomatic Medicine volume 47, No. 4, 1985). Further, providing intravenous testosterone to women as part of clinical studies is known (American Journal of Obstetrics and Gynecology, December 1986 pages 1288 to 1292).

15

#### SUMMARY OF THE INVENTION

Formulations for aerosolized delivery are disclosed. The formulations are preferably present in a container which can be easily loaded into an aerosol drug delivery device. The formulation is comprised of a testosterone and a suitable carrier which are combined to provide a formulation suitable for aerosol administration to a patient - preferably a human adult female.

The invention further provides kits which are comprised of a device for creating an aerosol and a plurality of containers adapted for being loaded into the device. The container holds a formulation comprised of testosterone and a carrier and is aerosolized to create particles having a diameter in a range of from about 1 micron to about 3 microns in diameter. Containers may hold a single dose or multiple doses and the active ingredient is present in a therapeutically effective amount. The ingredient is a testosterone and/or a vasodilator, e.g. sildenafil citrate.

A method of increasing the libido of a woman over a discrete period of time (e.g. 30-240 minutes) by the administration of testosterone is disclosed. A formulation comprised of testosterone is aerosolized preferably producing particles which have a size in a range of from about 1 to 3 microns which can be inhaled into areas of the lung where they can readily enter the blood stream. The aerosol is inhaled into the lungs of a patient. Once inhaled,

particles of testosterone deposits on lung tissue and from there enter the patient's circulatory system and thereby increase the patient's serum testosterone level. The percentage increase in the patient's testosterone level will vary depending on the needs of the patient. However, the patient's normal baseline serum testosterone level is preferably increased 25% or more and more preferably 100% or more. Because delivery is by inhalation the patient's serum testosterone level is quickly raised to a desired level e.g. in thirty minutes or less, more preferably fifteen minutes or less. When the patient's testosterone level is raised to a desired level the patient's libido is increased. The increased testosterone level gradually subsides (as the testosterone is metabolized and cleared) thereby avoiding the adverse side effects generated by maintaining enhanced testosterone levels over long periods.

An aspect of the invention is a method of treatment whereby testosterone or derivative thereof is aerosolized, inhaled and provided to the circulatory system of the patient at levels sufficient to increase libido (over a short period of time) and propensity for orgasm.

An advantage of the invention is that the testosterone levels are raised within minutes of administration and return to normal levels within hours -- preferably in less than four hours.

Another advantage is that the administered testosterone is quickly metabolized allowing the patient's testosterone levels to return to normal thereby avoiding the adverse effects of long term administration.

A feature of the invention is that aerosolized particles of testosterone having a diameter of about 0.5 to 8 microns (preferably 1-3 microns) are created and inhaled deeply into the lungs thereby enhancing the speed and efficiency of administration.

It is an object of this invention to describe the utility of delivering testosterone or dihydrotestosterone by inhalation as a means of treating women with decreased libido and/or decreased propensity to have orgasms.

It is another object of this invention to describe liquid formulations (which includes suspensions) of testosterone and derivatives thereof such as dihydrotestosterone appropriate for pulmonary delivery.

It is another object of this invention to describe how testosterone or dihydrotestosterone delivered via the lung can quickly increase plasma levels substantially beyond baseline levels for the patient.

It is another object of this invention to describe the blood levels of testosterone or dihydrotestosterone required for rapid onset of a normal to enhanced libido in women with baseline decreased libido.

5 It is another object of this invention to describe the time course of inhalation of testosterone or dihydrotestosterone and the onset of increased libido in women suffering from decreased libido.

It is another object of this invention to describe how the pulsatile delivery of testosterone or dihydrotestosterone as replacement therapy for women with decreased libido is associated with a decreased incidence of side effects (secondary sexual characteristics) commonly associated with traditional testosterone replacement therapy which produces a steady state level of the hormone.

The delivery of testosterone by inhalation provides, for the first time, the means for non-invasively delivering clinically relevant amounts of testosterone on demand near the time of planned intercourse.

15 It is an object of the invention to provide a method of treatment of erectile dysfunction in a patient comprising the steps of aerosolizing a formulation comprising sildenafil citrate, inhaling the aerosolized formulation into the lungs of a patient, and allowing the particles of sildenafil citrate to deposit on lung tissue and enter the patient's circulatory system.

20 It is an object of the invention to provide an aerosolized formulation comprised of sildenafil citrate and a carrier, the aerosol being characterized by particles having a diameter in the range of about 1.0 micron to 5.0 microns making up 50% or more of the aerosol particles.

It is an object of the invention to provide a kit comprising an aerosol delivery device and a formulation comprising a testosterone, sildenafil citrate, or a combination thereof.

25 It is an object of the invention to provide a kit comprising two aerosol delivery devices and two formulations, a first formulation comprising a testosterone for use by a women, and a second formulation comprising a testosterone, sildenafil citrate, or a combination thereof, for use by a man.

30 These and other aspects, objects, advantages, and features of the invention will become apparent to those skilled in the art upon reading this disclosure.

## DEFINITIONS

The terms "testosterone", "a testosterone" and the like are used interchangeably here and are intended to mean the naturally occurring hormone known as testosterone having the chemical name 17- $\beta$ -hydroxyandrost-4-en-3-one which may be isolated and purified from nature or synthetically produced in any manner. These terms are also intended to encompass the commonly occurring reduced version of testosterone having been reduced by 5 $\alpha$ -reductase to 5 $\alpha$ -dihydroxytestosterone which is also referred to here as dihydrotestosterone or simply "a testosterone." A dihydrotestosterone may be isolated from nature but is preferably synthetically produced and purified. Testosterone USP is a white or creamy-white crystalline powder having a molecular weight of 288.43.

The term "testosterone derivative" refers to any androgen hormone for pharmaceutical use. The term includes testosterone esters, i.e. compounds where the "H" of the "OH" group is replaced with an alkyl group, e.g. propionate, cypionate and enanthate. Other pharmaceutically acceptable derivatives include methyltestosterone, methandrostenedione, fluovymesterone and danazol. A number of useful derivatives of testosterone are disclosed within the Physician's Desk Reference (most recent edition) as well as Harrison's Principles of Internal Medicine. In addition, applicants refer to U.S. Patents 5,536,714 issued July 16, 1996; 5,824,668 issued October 20, 1998; 3,980,638 issued September 14, 1996; 4,031,117 issued June 21, 1977; 4,085,202 issued April 18, 1978; 4,197,286 issued April 8, 1980; 4,507,290 issued March 26, 1985 and 5,622,944 issued April 22, 1997 all of which are incorporated herein by reference to disclose and describe testosterone derivatives and formulations.

The terms "diameter", "particle diameter" and the like are used interchangeably herein to refer to particle size as given in the "aerodynamic" size of the particle. The aerodynamic diameter is a measurement of a particle of unit density that has the same terminal sedimentation velocity in air under normal atmospheric conditions as the particle in question. This is pointed out in that it is difficult to accurately measure the diameter of small particles using current technology and the shape of such small particles may be continually changing. Thus, the diameter of one particle of material of a given density will be said to have the same diameter as another particle of the same material if the two particles have the same terminal sedimentation velocity in air under the same conditions. In connection with the present invention it is important to have particles which do not have too large of a

diameter so that the particles can be inhaled deeply into the lungs and thereby deposited on lung tissue and transferred into the patient's circulatory system. It is equally important not to have particles which are too small in that such particles would be inhaled into the lungs and then exhaled without depositing on the lung tissue in the same manner that particles of smoke can be inhaled and exhaled with only a small amount of the particles being deposited on the lung tissue. An acceptable range for particle diameter is in the range of 0.5 to 12 microns, preferably 0.5 to 8 microns and more preferably 1 to 3 microns.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Before the devices, formulations, and methodology of the present invention are described, it is to be understood that this invention is not limited to the particular device, components, formulations and methodology described, as such may, of course, vary. It is also to be understood that the terminology used herein is with the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "and," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a formulation" includes mixtures of different formulations and reference to "the method of treatment" includes reference to equivalent steps and methods known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to describe and disclose specific information for which the reference was cited in connection with.

All publications mentioned herein are incorporated herein by reference to described and disclose specific information for which the reference was cited in connection with. The publications discussed herein are provided solely for their stated disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such publications by virtue of prior invention. Further,



the actual publication date may be different from that stated on the publication and as such may require independent verification of the actual publication dates.

### INVENTION IN GENERAL

5           Despite the fact that steady state delivery of testosterone as replacement therapy for women experiencing decreased libido is inherently prone to producing unwanted side effects, the use of pulsatile testosterone replacement therapy to mimic the normal elaboration of this hormone during ovulation has not been explored. The use of testosterone replacement therapy for brief courses of treatment has been attempted, however the slow rate  
10 of absorption of methyl testosterone from pills has limited its utility. In order to replace the missing testosterone in a therapeutically effective manner, it is necessary to provide a rapid pulse of bioavailable testosterone to the patient on demand. In this way, testosterone could be replaced by the patient as needed coincident with the desire to engage in sexual activity.

          It is not surprising that clinical studies evaluating the effect of acute, on demand  
15 testosterone replacement therapy in women with decreased libido have not been attempted. The only tool currently available for a true pulsatile, rapid onset replacement therapy is intravenous administration. Although preparations of testosterone appropriate for intravenous administrations have been available for some time, intravenous cannulation as the means for gaining access to the circulation for the administration of testosterone on  
20 demand is inconsistent with the desire for women to be able to modulate their libido in concert with the course of their daily lives.

          Precision delivery of small molecule drugs via the lung for systemic effect is possible. An electronic inhaler capable of delivering a liquid formulated drug stored in a unit dose packages has been described. A formulation of testosterone or dihydrotestosterone  
25 can be prepared for delivery with this system. The quantitative delivery of testosterone or dihydrotestosterone, on demand by a woman prior to initiation of sexual intercourse, provides a mechanism for testosterone replacement therapy which is unlikely to be associated with side effects precipitated by chronic delivery of the drug.

          While particularly applicable to post menopausal woman, the use of testosterone  
30 replacement therapy to modulate libido could be of value to women still of child bearing age. Disappearance of or reduction of the libido has been described in women who are continuing to ovulate. The reduction in libido may be due to therapy including the use of birth control

pills which contain hormones. Therefore, acute administration of testosterone to significantly raise blood levels for discrete periods has potentially widespread application in women across a wide range of ages.

The baseline serum testosterone level of a normal adult human female is generally below about 1 ng/ml with modest changes through the menstrual cycle (Geobelman et al., *Am J. Obstet. Gynecol.* 119:445 (1974)) with general fluctuation between about 0.3 to 0.5 ng/ml. However, adult human females with polycystic ovarian disease have ovarian vein testosterone levels of 20 to 65 ng/ml and peripheral venous levels of about 7.5 ng/ml (Dupon et al., *Am. J. Obstet. Gynecol.* 115:478 (1973)). Abnormally high levels of testosterone over long periods are associated with acne and hirsutism.

To maintain normal testosterone levels an adult human female will produce about 0.25 mg of testosterone per day as compared to about 5-6 mg/day produced by a normal adult male to maintain a normal adult male testosterone level of 3 to 10 ng/ml. Because women produce such small amounts of testosterone the administration of very small amounts will dramatically increase the patient's normal levels. In accordance with the present invention 0.05 mg to 5 mg, preferably 0.25 to 2 mg and more preferably about 1 mg of testosterone is administered to the circulatory system of the patient. Administration of such amounts to the circulatory system may require aerosolizing larger amounts due to inefficiencies in the aerosol delivery system.

Testosterone can be administered orally. However, after oral administration it is absorbed from the gut into the portal blood and degraded promptly by the liver. Thus, insignificant amounts reach the patient's systemic circulation. Testosterone can also be administered parenterally. However, when so administered it is rapidly absorbed and metabolized making it difficult to sustain effective levels in plasma over time. In view of such, effective therapy has been carried out using means of delivery where testosterone is slowly absorbed (e.g. dermal patches) or when the testosterone is chemically modified to retard absorption and/or catabolism.

The present invention uses intrapulmonary delivery to avoid first pass liver metabolism and to obtain quick infusion into the patient's systemic circulatory system. Further, the method of the present invention does not require maintaining increased testosterone levels over long periods. Accordingly, chemical modification to retard absorption and/or catabolism are not required or desired.

The present invention administers sufficient testosterone by inhalation to temporarily raise the patient's libido, increase the patient's propensity for orgasm, and thereafter allow the patient's testosterone level to return to a level normally experienced by the patient. Because intrapulmonary administration is not 100% efficient the amount of drug aerosolized will be greater than the amount which actually reaches the patient's circulation. For example, if the inhalation system used is only 50% efficient then the patient will aerosolize a dose which is twice that needed to raise the patient's testosterone level to the extent needed to obtain the desired results. More specifically, when attempting to administer 1 mg of testosterone with a delivery system known to be 50% efficient the patient will aerosolize an amount of formulation containing about 2 mg of testosterone.

Testosterone therapy as described herein can be used in construction with other therapies intended to increase or enhance libido. Such therapies include but are not limited to herbal preparations and vitamin supplements.

## INDICATIONS

The method of the invention has broad applicability to both the male and female populations. However, its use is specifically indicated in four categories.

First, post-menopausal women who have experienced all or any of (1) decreased levels of testosterone; (2) decreased libido; and (3) decreased propensity to experience orgasm.

Second, women of child bearing age who have experienced all or any of (1) decreased levels of testosterone; (2) decreased libido; and (3) decreased propensity to experience orgasm.

Thirdly, women of child bearing age being treated with birth control pills who have experienced all or any of (1) increased levels of estrogen relative to testosterone resulting in either or both of (2) decreased libido and (3) decreased propensity to experience orgasm.

Fourthly, men having erectile dysfunction symptoms, especially those due to peripheral vascular disease.

Fifthly, men having a decreased level of serum testosterone.

In the first three categories it is not desirable to administer sufficient amounts of a testosterone so as to raise the patient's testosterone level continually over long periods of time. For example, it is not desirable to administer testosterone several times per day for

several days. Such will raise testosterone levels over long periods and result in adverse side effects including acne, and increased growth of body hair.

## DOSING

5           The amount of a testosterone administered will vary based on a factor such as the  
age, weight and baseline testosterone level of the patient. Initially, small doses, e.g. about  
0.25 mg, is administered. If the desired result is obtained no further dosing is provided. If  
the desired effect is not obtained additional 0.25 mg doses can be administered up to 2.0 mg.  
If the patient finds that larger doses are needed then for further treatment the patient may be  
10       provided with doses of 0.5 mg, 1.0 mg or 2.0 mg. The amount aerosolized may be  
substantially greater than the amount administered if the interpulmonary delivery device is  
inefficient. Thus, the device and method efficiencies must be taken into consideration when  
calculating the doses.

When testosterone enters the circulatory system of a human patient it is readily  
15 reduced via 5 $\alpha$ -reductase to 5 $\alpha$ -dihydrotestosterone. Thus, when referring to increasing a  
patient's testosterone level this disclosure is referring to combined levels of testosterone and  
5 $\alpha$ -dihydroxytestosterone present in the patient's serum. The present invention includes the  
administration of 5 $\alpha$ -dihydroxytestosterone which is the active molecule. The invention also  
includes the administration of testosterone derivatives provided such derivatives increase  
20 libido and do not result in unacceptable adverse effects.

Obtaining a result such as increased libido may be difficult to ascertain. Some placebo effect will be experienced by some patients and others may continuously administer doses in an attempt to obtain a more enhanced effect. To avoid undesirable side effects from overdosing or from dosing too frequently the delivery device may be controlled by a suitable lockout system such as taught in U.S. Patents 5,507,277; 5,694,919; and 5,735,263. Such a system can prevent release of more than a given amount of drug at a single dosing event and/or restrict the number of dosing events within a given period of time. The restrictions are designed to prevent the patient from experiencing adverse secondary effects.

## 30 FORMULATIONS/DEVICES

Pharmaceutical grade testosterone can be produced as a white or creamy white powder. The pure powder is aerosolized and inhaled by itself or with the use of a dry

powder inhaler (DPI) device. However, it is desirable to formulate the crystals with an excipient to provide small particles of dry powder which do not stick together. The particles preferably have a diameter in a range of from about 1 to 10 microns more preferably 1 to 5 microns and still more preferably about 1 to about 3 microns. Methods of formulating dry  
5 powders and dry powder inhaler devices are disclosed in U.S. Patents 5,826,633; 5,814,607; 5,785,049; 5,780,014; 5,775,320; 5,740,794; and Des. 390,651 all of which are incorporated by reference to describe and disclose such.

Testosterone is relatively insoluble in water. Accordingly, to create a solution of testosterone a solubilizer (e.g., one of several cyclodextrines or phospholipids) or an organic  
10 solvent such as ethanol is used. Alternatively, a microsuspension of testosterone in water with or without ethanol can also be produced. The testosterone solution is aerosolized and inhaled. The testosterone solution can be placed in a low boiling point propellant in a pressurized canister and released using a conventional metered dose inhaler (MDI) device. Preferably, the MDI device is modified so that the aerosolized dose is released each time at  
15 the same inspiratory flow rate and inspiratory volume. When this is done the patient is more likely to receive the same dose each time. A device for obtaining repeatable dosing with an MDI canister is taught in U.S. Patent 5,404,871 issued April 11, 1995.

In accordance with the present invention it is preferable to load the testosterone solution into a container which opens to a porous membrane. When the formulation is  
20 moved through the membrane it is aerosolized. Such containers are taught in U.S. Patent 5,497,763 issued March 12, 1996. The container is loaded into a device and delivered via a method as taught in U.S. Patent 5,823,178 issued October 10, 1998 both of which patents are incorporated herein by reference to describe and disclose containers, devices and methods of drug delivery by inhalation.

25 Aerosol drug delivery devices vary but generally are comprised of (1) a container for the drug e.g. testosterone; (2) a means for aerosolizing the drug; and (3) a mouthpiece from which the aerosol is inhaled. The aerosol can be any small particles dispersed in air, e.g. a cloud of a dry powder or a fine spray of liquid formulation. Nebulizers, metered dose inhalers (MDIs) and dry powder inhalers (DPIs) are the most well known devices for  
30 creating an aerosol. Less conventional devices known as electrohydrodynamic aerosol devices as taught in U.S. Patent 4,358,059; PCT WO 99/07478; and 98/03267 can also be used to create an aerosol in the method of the invention.

Nasal or buccal formulations could be used for nasal or buccal delivery.

#### AEROSOL ADMINISTRATION OF SILDENAFIL

Similarly, agents such as sildenafil citrate (U.S. 5,426,107 and U.S. 5,250,534) can  
5 also be administered to women or men by the methods of the instant invention, alone or in  
combination with testosterone.

Sildenafil citrate, also termed VIAGRA™, is typically administered in tablet form to  
men experiencing erectile dysfunctions resulting from peripheral vascular disease. The  
tablet is taken orally about thirty (30) minutes to four (4) hours before sexual activity.

10 The typical oral dose of sildenafil is 25-100mg per day. As described above, such  
doses can be administered to the lungs through the use of aerosolized aqueous solutions or as  
a dry powder.

The sildenafil citrate can be formulated alone or as an admixture with a testosterone  
for simultaneous delivery.

15 In an embodiment, men in need of supplemental testosterone and experiencing  
erectile dysfunction due to peripheral vascular diseases can administer the drugs either  
simultaneously or sequentially. The dosage of testosterone will typically be sufficient to  
raise the serum testosterone level of the man to a normal range, that being about 200 - 1000  
ng/dL.

20 The formulation of sildenafil citrate can be administered to male or female patients  
and may be administered alone or in combination with an aerosolized dose of testosterone.

In one embodiment of the invention the sildenafil citrate is administered orally and  
the testosterone is administered by aerosol about 30 - 60 minutes after the oral  
administration of sildenafil citrate. The oral administration of sildenafil citrate is an  
25 administration in advance of a sexual event and after allowing time to achieve a therapeutic  
effect on increasing blood flow the patient is dosed with testosterone via aerosol. The  
testosterone enhances the libido and the sildenafil citrate enhances the patient's ability to  
perform and/or achieve orgasm.

The aerosolized administration of testosterone could also be in combination with  
30 other drugs used in the treatment of various sexual dysfunctions e.g. administered in  
combination with the topical application of alprostadil. Other oral, injectable and topical  
drugs are and will become available for the treatment of sexual dysfunctions and such drugs

(e.g. vasodilators) can be used in combination with aerosolized delivery of testosterone to obtain enhanced results. It is noted that although such drugs may, by themselves, facilitate sexual activity they do not effect libido. Accordingly, a truly enhanced effect is obtainable by combining a drug which increases blood flow to a desired area with aerosolized delivery of testosterone which increases libido.

### KITS

In an embodiment of the invention, a kit is provided for use by a healthcare provider, more preferably a patient. An exemplary kit will provide a hand-held aerosol delivery device and at least one dose, preferably one to about one hundred, more preferably one to thirty doses of a testosterone for use by a woman. In an embodiment, the kit will comprise a hand-held aerosol delivery device and at least one dose, preferably one to about one hundred, more preferably one to thirty doses of a testosterone for use by a man. In an embodiment, the kit will provide a hand-held aerosol delivery device and at least one dose, preferably one to about one hundred, more preferably one to thirty doses of an admixture of testosterone and sildenafil citrate for use by a man. In an embodiment, the kit will contain a hand-held aerosol delivery device and at least one dose, preferably one to about one hundred, more preferably about one to thirty doses of sildenafil citrate for use by a man.

In an embodiment, a kit is provided which comprises two hand-held delivery devices, wherein a first delivery device comprises at least one dose, preferably one to one hundred doses, of a testosterone for use by a woman. The second delivery device comprises at least one dose, preferably one to one hundred doses, of a testosterone, sildenafil citrate, or a combination thereof for use by a man. Such a kit is intended for use by a couple in need of such treatment.

The kit of the invention can be comprised of various combinations of drugs and drug delivery devices. However, the kit will preferably be comprised of an aerosol drug delivery device which comprises a container which holds one or a plurality of doses of testosterone, a means for aerosolizing the testosterone and a mouthpiece from which the aerosolized testosterone may be inhaled. This device is present in the kit with another drug. For example, the kit may comprise a container of sildenafil citrate or related drug which obtains a response similar to sildenafil citrate. The other drug may be administered orally or topically but is preferably in a container which can be loaded into the device used to deliver

the testosterone by inhalation. Thus, a preferred kit will comprise a drug delivery device which can generate an aerosol for inhalation and a plurality of containers of testosterone which can be loaded into the device and a plurality of containers of a vasodilator such as sildenafil citrate which can be loaded into the device.

- 5           The instant invention is shown and described herein in a manner which is considered to be the most practical and preferred embodiments. It is recognized, however, that departures may be made therefrom which are within the scope of the invention and that obvious modifications will occur to one skilled in the art upon reading this disclosure.



What is claimed is:

1. A formulation for aerosolized administration of a testosterone, comprising:  
a testosterone; and  
5 a carrier suitable for aerosolized delivery.
2. The formulation of claim 1, characterized by being non-irritating to a patient's respiratory tract.
- 10 3. The formulation of claim 2, wherein the formulation is in a container suitable for loading into an aerosol drug delivery device.
4. The formulation of claim 3, wherein the container holds an amount of testosterone such that when aerosolized and administered to a patient the patient's  
15 testosterone level is increased from a first level to a second increased level which is twice or more the first level in a period of time of thirty minutes or less.
5. The formulation of claim 4, wherein the aerosolized formulation comprises 0.25 mg or more of testosterone and further wherein the patient's testosterone level is  
20 increased to 0.8 ng/ml or more.
6. The formulation of claim 1, wherein the testosterone is dihydrotestosterone, and the patient's dihydrotestosterone level is increased to about 0.8 ng/ml in fifteen minutes or less.
- 25 7. The formulation of claim 6, further comprising sildenafil citrate.
8. The formulation of claim 1, wherein the formulation is a dry powder formulated for inhalation from a dry powder inhaler device.
- 30 9. A formulation for increasing libido by aerosol administration, comprising:  
a testosterone; and

a carrier suitable for aerosolized administration.

10. The formulation of claim 9, wherein the testosterone is dihydrotestosterone.
- 5 11. The formulation of claim 9, wherein aerosolization of the formulation provides aerosolized particles having a diameter in a range of from about 1 to about 5 microns.
12. The formulation of claim 9, wherein the formulation is a dry powder
- 10 formulation.
13. The formulation of claim 9, wherein the formulation is a liquid formulation and the carrier is selected from the group consisting of an organic solvent to form solutions and a liquid to form suspensions.
- 15 14. An aerosolized formulation comprised of a testosterone and a carrier, the aerosol being characterized by particles having a diameter in a range of about 1.0 micron to 5.0 microns making up 50% or more of the aerosol particles.
- 20 15. An aerosolized formulation comprising a pharmaceutically acceptable salt of sildenafil and a carrier, the aerosol being characterized by particles having a diameter in a range of about 1.0 micron to 5.0 microns making up 50% or more of the aerosol particles.
- 25 16. A kit, comprising:  
an aerosol delivery device; and  
a formulation comprising an active ingredient selected from the group consisting of testosterone, sildenafil citrate, a combination thereof.

17. The kit of claim 16, comprising:

· a plurality of containers of testosterone which containers are adapted for use with the aerosol delivery device; and

5 a plurality of containers of sildenafil citrate which containers are adapted for use with the aerosol delivery device.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/12092

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 9/12

US CL : 424/45

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/45

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,426,107 A (BELL et al.) 20 June 1995, see entire document.	1-17
Y	US 5,536,714 A (KOJIMA et al.) 16 June 1996, see entire document.	1-17



Further documents are listed in the continuation of Box C.



See patent family annex.

* "A" document defining the general state of the art which is not considered to be of particular relevance	** later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"B" earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"Z" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 JULY 2000

Date of mailing of the international search report

18 AUG 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAJ BAWA, PH.D.

Telephone No. (703) 308-1235

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number  
WO 00/66206 A3

- (51) International Patent Classification: A61M 15/02, A61K 9/72
- (21) International Application Number: PCT/US00/11799
- (22) International Filing Date: 2 May 2000 (02.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/132,215 3 May 1999 (03.05.1999) US  
Not furnished 3 February 2000 (03.02.2000) US
- (71) Applicant: BATTELLE MEMORIAL INSTITUTE  
[US/US]; 505 KinG Avenue, Columbus, OH 43201-2693 (US).
- (72) Inventors: THURSTON, Rachel, M.; 7768 Heathermoon Drive, Columbus, OH 43235 (US). BROWNING, James, D.; 198 Richards Road, Columbus, OH 43214 (US). SHAH, Frafal, K.; 4750 Coolbrook Drive, Columbus, OH 43026 (US). PLACKE, Michael, E.; 1473 Inglis Avenue, Columbus, OH 43212 (US).
- (54) Title: COMPOSITIONS FOR AEROSOLIZATION AND INHALATION
- (57) Abstract: A composition used in combination with an electrohydrodynamic device capable of delivering an active ingredient to the aerodigestive system of the user. The composition comprises three or optionally four basic components: an active ingredient; a carrier material in which the active ingredient may be dissolved, suspended, or emulsified; an aerosol properties adjusting material which provides the composition with the physical characteristics required to create an aerosol cloud by electrostatic or electrohydrodynamic means; and optionally at least one excipient that further adjusts, preserves, stabilizes, or enhances the overall performance of the composition.
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— With international search report.
- (88) Date of publication of the international search report:  
8 February 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/66206 A3

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/US 00/11799

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61M15/02 A61K9/72

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61M A61K B05B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 234 842 A (IMPERIAL CHEMICAL INDUSTRIES PLC, UK) 2 September 1987 (1987-09-02) claims 1-3,7 page 7, line 32 - page 8, line 18 page 14, line 27 - page 16, line 6	1, 2, 6, 15, 17-22
X	WO 95 26235 A (ELECTROSOLS LTD., UK) 5 October 1995 (1995-10-05) claims 1, 5-7 page 2, line 19 - page 3, line 5 page 3, line 21 - line 33 page 5, line 10 - line 23	1, 2, 15, 17, 18, 21, 22

-/-



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*S\* document member of the same patent family

Date of the actual completion of the international search

31 October 2000

Date of mailing of the international search report

07/11/2000

Name and mailing address of the ISA  
European Patent Office, P.B. 5618 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Scarponi, U

# INTERNATIONAL SEARCH REPORT

Int. Jonal Application No  
PCT/US 00/11799

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 99 49981 A (CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE,FR) 7 October 1999 (1999-10-07) claims 1-3,11-15 page 15, line 16 - line 24 page 16, line 7 - line 8 page 20, line 15 -page 21, line 5 page 24, line 3 - line 5 -----	1-3, 6, 15,17-22
X,P	WO 99 42153 A (BESPAK PLC,UK) 26 August 1999 (1999-08-26) claims 1,9,11 page 4, line 25 - line 35 -----	1,2,15, 17-22
E	WO 00 38770 A (BATTELLE MEMORIAL INSTITUTE,U.S.A.) 6 July 2000 (2000-07-06) claims 1,4,5,7,11,21,22,27,28,39,44,45,48,50-52 page 1, line 6 - line 12 page 4, line 10 page 5, line 18 - line 21 page 7, line 22 -page 8, line 4 page 15, line 1 - line 23 page 27, line 5 - line 6 -----	1,2,15, 17-22
E	WO 00 35524 A (ELECTROSOLS LTD.,UK) 22 June 2000 (2000-06-22) claims 1,2,20,22,26,34,40-44,46-56 page 3, line 10 - line 18 page 4, line 5 - line 9 page 4, line 16 -page 5, line 7 page 24, line 12 - line 17 page 42, line 17 -page 43, line 11 page 43, line 24 - line 27 page 44, line 16 - line 21 page 48, line 1 - line 3 tables 1,2 -----	1,2,8, 10,17-22

## INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. l. Application No

PCT/US 00/11799

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 234842 A	02-09-1987	AT 56156 T	15-09-1990
		AU 582949 B	13-04-1989
		AU 6906187 A	27-08-1987
		AU 594429 B	08-03-1990
		AU 6906287 A	27-08-1987
		CA 1275883 A	06-11-1990
		DE 3764662 D	11-10-1990
		DK 82587 A, B,	22-08-1987
		EP 0234841 A	02-09-1987
		FI 870639 A, B,	22-08-1987
		FI 870640 A	22-08-1987
		GR 3000843 T	15-11-1991
		IE 59596 B	09-03-1994
		IL 81572 A	15-01-1992
		JP 2109019 C	21-11-1996
		JP 8024714 B	13-03-1996
		JP 62197071 A	31-08-1987
		JP 62254830 A	06-11-1987
		NO 870658 A, B,	24-08-1987
		NO 870660 A	24-08-1987
		NZ 219305 A	26-02-1990
		NZ 219306 A	26-02-1990
		PT 84320 A, B	01-03-1987
		US 4829996 A	16-05-1989
		US 4795330 A	03-01-1989
		ZA 8701146 A	28-10-1987
		ZA 8701147 A	25-11-1987
WO 9526235 A	05-10-1995	DE 69510046 D	08-07-1999
		DE 69510046 T	23-09-1999
		EP 0752918 A	15-01-1997
		JP 9510653 T	28-10-1997
		US 5813614 A	29-09-1998
WO 9949981 A	07-10-1999	FR 2776538 A	01-10-1999
		AU 2940599 A	18-10-1999
WO 9942153 A	26-08-1999	GB 2334461 A	25-08-1999
		AU 1163999 A	06-09-1999
WO 0038770 A	06-07-2000	AU 2592900 A	31-07-2000
WO 0035524 A	22-06-2000	GB 2345010 A	28-06-2000
		AU 1872500 A	03-07-2000